



Tropentag, October 9-11, 2007, Witzenhausen

“Utilisation of diversity in land use systems:  
Sustainable and organic approaches to meet human needs”

**Potential of the Parasitic Wasp, *Lariophagus distinguendus* (Förster) (Hymenoptera: Pteromalidae) as a Biological Control Agent for *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae) in Stored Maize**

CHARLES ADARKWAH<sup>1,4</sup>, DANIEL OBENG-OFORI<sup>2</sup>, SABINE PROZELL<sup>3</sup>, MATTHIAS SCHÖLLER<sup>3</sup>,  
CHRISTOPH REICHMUTH<sup>4</sup>, CARMEN BUETTNER<sup>1</sup>

<sup>1</sup>Humboldt-Universität zu Berlin, Institute for Horticultural Sciences, Section Phytomedicine, Germany

<sup>2</sup>University of Ghana, Crop Science, Ghana

<sup>3</sup>Biological Consultants Ltd., Germany

<sup>4</sup>Federal Biological Research Centre for Agriculture and Forestry (BBA), Institute for Stored Product Protection, Germany

**Abstract**

The parasitic wasp, *Lariophagus distinguendus* is an ectoparasitoid of several beetle species that feed on durable stored products. The potential using *L. distinguendus* for the biological control of *S. zeamais* was assessed in maize stored in storage cylinders. The host finding behaviour of the parasitoid was studied in maize stored in various vertical depths in the cylinders. Holes of 3 mm diameter were drilled through PVC pipes of 20.5 cm length and 20 mm diameter. The pipes were inserted into the holes in the cylinders. An acoustic detector served to identify the maize kernels that contained 3 weeks old larvae of *S. zeamais*. Uninfested maize kernels were filled into the cylinder to depths of 20, 25, 30, 35, 40, 45, 95 and 100 cm, respectively. For depths of 20, 25 and 30 cm, 25 adult *L. distinguendus* aged between 0–14 days were released; for 35, 40 and 45 cm, 30 adult *L. distinguendus* were released while for 95 and 100 cm, 100 adult *L. distinguendus* of the same age were released, each on top of the uninfested maize. Each treatment was repeated three times with control without parasitoids. *L. distinguendus* adults that entered the pipe and the wire mesh cage to parasitise the *S. zeamais* infested maize kernel were collected and placed in a 250 ml glass jars. The emergence of *S. zeamais* was recorded in both *L. distinguendus* treated and untreated maize weekly until the 6<sup>th</sup> week. *L. distinguendus* penetrated and infested *S. zeamais* stored in the cylinders at the various depths. This showed that *L. distinguendus* was able to find its host in the cylinder with infested maize kernels and produced F1 offspring. *L. distinguendus* also significantly reduced the emergence of *S. zeamais* in stored maize. The significance of these results with respect to the suitability of *L. distinguendus* for the biological control *S. zeamais* is discussed.

**Keywords:** Maize, parasitic wasp, storage