

Ecological trade-offs between plant biodiversity, land-uses and management intensification: study on four contrasting agroforestry land-uses in tropical forests of Bangladesh

Sharif Ahmed Mukul^{1,2*}, Narayan Saha³

¹*Faculty of Life Sciences, Centre for Forests, Landscape and Planning, Copenhagen University, Copenhagen, Denmark;*

²*Centre for Research on Land-use Sustainability, Noakhali 3800, Bangladesh;*

³*Department of Forestry and Environmental Science, School of Agriculture and Mineral Sciences, Shahjalal University of Science and Technology, Sylhet 3114, Bangladesh*

**Corresponding author; e-mail: sharif_a_mukul@yahoo.com*

Abstract

Agroforestry, because of its diverse options and structure sometimes believed to more suitable for conserving biodiversity, particularly in tropical landscapes where rural people depends heavily on nature for sustaining their livelihoods. However, since agroforestry systems are subjected to some level of cultural management it is also critical to understand how cultural managements and different level of intensification affect biodiversity in agroforestry landscapes. We conducted an exploratory survey on four contrasting agroforestry systems; viz., betel-vine (*Piper betel*) based agroforestry, lemon (*Citrus limon*) based agroforestry, pineapple (*Ananas comosus*) based agroforestry, and short-rotation shifting cultivation in a tropical forest patch of Bangladesh, to assess the response of plant community to different level of management intensification. A unique management intensification gradient was created identifying a total of 11 locally used cultural practices followed across the studied agroforestry systems. We recorded 173 plant species (61 trees, 42 shrubs, 47 herbs, 18 climbers and 5 orchids) from forty 100m² plots established in four agroforestry land-uses. The Shannon-Weiner biodiversity index calculated was higher for betel-vine agroforestry (3.3), followed by 2.9 for lemon/horticulture agroforestry, 2.2 for pineapple agroforestry and 1.9 in short-rotation shifting cultivation system (comparable to 2.7 for forest). Species evenness index for tree was also higher in betel-vine agroforestry system (0.79). A simple linear regression was performed for each of the studied agroforestry systems to realize the response of different plant functional groups to a weighted management intensification value. It was observed that, plant biodiversity is highly sensitive to management intensification, and the response of different plant functional groups to different management regimes varies greatly on the type of agroforestry systems and the kind of expected products. Study also suggested that, some practice (for example here betel-vine agroforestry system) that evolved through indigenous innovation is suitable for conservation of plant biodiversity and could offer a basis for sustainable forest management.

Key-words: agroforestry, management intensification, land-use, Lawachara National Park, Bangladesh
