



Tropentag 2014, Prague, Czech Republic
September 17-19, 2014

Conference on International Research on Food Security, Natural Resource
Management and Rural Development
organised by the Czech University of Life Sciences Prague

Effect of dietary sodium diformate in broilers on the productivity index against a positive control

Christian Lückstädt¹, Nicolas Greiffenstein² and Ronnie Dari³

¹ADDCON GmbH, Bonn, Germany; christian.lueckstaedt@addcon.com

²ADDCON GmbH, Bogota, Colombia

³Salus, Jaguariúna, Brazil

Abstract

Gastrointestinal diseases pose a serious threat to commercial poultry production. Organic acids and their salts are commonly used to suppress gastrointestinal disorders and thereby improve productivity. In a new commercial study, sodium diformate (NDF) was tested against a positive control (AGP) in order to investigate similar effects in poultry. The trial was conducted under large scale conditions in Ceará, Brazil in 2013 and aimed to test NDF against a commercial broiler diet containing an antibiotic growth promoter (Colistin). Around 500,000 one day old birds from 33 different houses were included in the study and were compared to the same number of birds (from the same houses) and their subsequent performance from the previous year. Performance data were measured at the end of the trial and the European Broiler Index (EBI) was calculated. The average growth period was 45 days. The positive control group used 60 g Colistin per t of feed for the first 21 days, while the treatment group added 0.1% NDF during the same time period. EBI in the houses with 0.1% NDF was increased by 5.1% ($P=0.006$). Due to the inclusion of sodium diformate, EBI in 10 out of 33 houses reached a value above 300 (max. 331), whereas the Colistin-treated houses attained only a maximum EBI of 297. These findings lead to the conclusion that the addition of 0.1% sodium diformate considerably improves overall broiler performance, combining effects on daily gain, survival and feed efficiency, even when compared to the use of an AGP.

Introduction

Enteric disorders are one of the most important groups of diseases which affect poultry production world-wide and they are continuing to cause high economic losses due to increased mortality rates, decreased weight gain, increased medication costs, and increased feed conversion rates (HAFEZ, 2011). Estimates of the annual costs of poultry-associated cases of salmonellosis and campylobacteriosis, two major causes of foodborne diseases in humans, in the United States alone ranged from \$64 million to \$114.5 million and \$362 million to \$699 million, respectively (BRYAN AND DOYLE, 1995). In the past this hazard was controlled by the prophylactic use of antibiotics. Since the EU ban on antibiotic growth promoters (AGP) in 2006, which led other markets to consider similar steps, new strategies against gastrointestinal diseases in livestock have been developed. Organic acids and their salts are commonly used to suppress gastrointestinal disorders and thereby improve productivity. Potassium diformate for instance, is the first substance with EU-approval as a non-antibiotic growth promoter for pigs.

Improving broiler performance or hygienic conditions with the aid of organic acids has been reported by many sources, as reviewed by DESAI ET AL. (2007). An important limitation, however, is that organic acids are rapidly metabolised in the fore-gut (crop to gizzard) of birds, which will reduce their impact on growth performance. A new molecule (sodium diformate, NDF – see Figure 1) has been proven to be effective against pathogenic bacteria, including Salmonella, along the whole gastro-intestinal tract (LÜCKSTÄDT AND THEOBALD, 2009). The reduced impact of pathogenic bacteria on the broiler, as well as the improved gut microflora, leading to a state of eubiosis in treated chickens, suggests that including diformate in broiler diets will also result in improved bird performance. Several trials have also been carried out over the last half-decade world-wide that document positive effects on broiler performance.




Figure 1: Chemical structure of sodium diformate (NDF), consisting of formic acid (upper molecule) and sodium formate (lower molecule) combined with a hydrogen bond

A recent study (LÜCKSTÄDT, 2013) analyzed the average impact of dietary NDF from all studies on the effect on weight gain, feed efficiency, mortality and productivity, against a negative control. The final data-set contained the results of 8 documented studies, comprising 17 trials with NDF-inclusion, which ranged from 0.1% to 0.6%. Those studies were carried out between 2006 and 2012 in China, France, Russia, Taiwan, Thailand and Vietnam under both commercial and institutional conditions and included more than 36,700 broilers from different breeds (Arbor Acres, Cobb, Hubbard) raised to between 35 and 44 days. The average level of dietary NDF from the data-set in all treated broilers was 0.28%. The typical dosage for NDF in broilers under commercial conditions ranged from 1-3 kg/t feed, depending on age (dietary protein level) and hygienic status of the farm. As shown in Table 1, NDF inclusion resulted in a numerical increase in feed intake of 1.1% (P=0.22).

Table 1: Performance analysis of 17 trials with broilers, fed diets with NDF, expressed as an average percentage difference from negative control

Dosage	Feed intake	Weight gain	FCR	Survival	EBI*
0.28	+1.1	+5.2	-4.1	+2.3	+12.4
P-value	0.22	0.0001	0.002	0.034	0.0005

*EBI = European Broiler Index

However, although feed intake was not improved significantly by NDF inclusion, the performance of broilers, based on daily gain, was significantly increased by 5.2% (P<0.001). Furthermore, the FCR was also significantly improved (4.1%; P<0.01). Survival was increased on average by 2.3% (P<0.05). Finally, the European Broiler Index (EBI) improved highly

significantly by 12.4% ($P < 0.001$), due to the inclusion of NDF. This finding is a result of improvements in all 3 of these parameters, indicating a potential for greater economic output. In a new commercial study, NDF was tested against an antibiotic growth promoter (Colistin) in order to demonstrate similar effects in poultry where the use of AGP is still allowed, as in South America.

Material and Methods

The trial was conducted under large scale conditions in Ceará, Brazil in 2013 and aimed to test NDF against a commercial broiler diet containing an antibiotic growth promoter (Colistin). Feed and water were available *ad libitum*. Around 500,000 one day old birds from 33 different houses were included in the study and were compared to the same number of birds (from the same houses) and their subsequent performance from the previous year. Performance data were measured at the end of the trial and the European Broiler Index (EBI), which is assessed on the basis of daily gain, survival rate and feed efficiency - thus combining the three most important parameters in broiler production - was calculated:

$$\text{EBI} = \text{ADG [g]} \times \text{Survival [\%]} / (10 \times \text{FCR})$$

The average growth period was 45 days. The positive control group used 60 g Colistin per t of feed for the first 21 days, while the treatment group added 0.1% NDF during the same time period. EBI data were analysed using the t-test. The results are given as mean \pm SD with a confidence level of 95%.

Results and Discussion

EBI in the houses with 0.1% NDF was increased by 5.1% ($P = 0.006$) – Table 2. Due to the inclusion of sodium diformate, in 10 out of 33 houses EBI reached a value above 300 (max. 331), whereas the Colistin-treated houses attained only a maximum EBI of 297 (see also Figure 2).

Table 2: Productivity index of broilers fed with or without sodium diformate (NDF)

	Positive control (60 g/t)	0.1% NDF	P-level
EBI	266 \pm 15	279 \pm 27	0.006
max. EBI	297	331	-
min. EBI	232	237	-
EBI > 300 [n]	0	10	-

In broilers, improved zootechnical performance is thought to stem from both improvements in the intestinal microflora, as a result of suppressing pathogenic bacterial species; and improved protein digestion. GANGULY (2013) reported in this regard "...Organic acids reduce the growth of many pathogenic and non-pathogenic intestinal bacteria, decrease intestinal colonisation and reduce infections, ultimately decreasing inflammatory processes at the intestinal mucosa." Furthermore, VAN DER SLUIS (2000) estimated that poor intestinal integrity can result in economic losses of up to 10 cents per bird. As often seen with other additives, hygiene also plays some role in the performance achieved. The average impact of NDF against an antibiotic growth promoter on the performance index remained above that normally expected. It can therefore be speculated that the benefit to feed efficiency found when NDF was included in diets, may not only stem from better gut integrity, but also as a direct result of the additive's impact on protein

digestibility. This was suggested, for instance, by SELLE ET AL. (2004) who found that the use of diformates in broilers improved nitrogen retention.




Figure 2: EBI on farms in eastern Brazil in broilers fed with NDF or Colistin (PC)

Conclusions and Outlook

These findings lead to the conclusion that the addition of 0.1% sodium diformate considerably improves overall broiler performance, combining effects on daily gain, survival and feed efficiency, even when compared to the use of an AGP.

References

- BRYAN, F.L. AND DOYLE, M.P. (1995). Health risks and consequences of *Salmonella* and *Campylobacter jejuni* in raw poultry. *Journal of Food Protection* 3: 229-344.
- DESAI, D., PATWARDHAN, D. AND RANADE, A. (2007). Acidifiers in Poultry Diets and Poultry Production. In: LÜCKSTÄDT, C.: *Acidifiers in Animal Nutrition – A Guide for Feed preservation and Acidification to Promote Animal Performance*. Nottingham University Press, Nottingham, 63-69.
- GANGULY, S. (2013). Potential non-antibiotic growth promoting dietary supplements for animal nutrition: A Review. *Journal of Applied Pharmaceutical Science* 3(7):174-178.
- HAFEZ, H.M. (2011). Enteric diseases of poultry with special attention to *Clostridium perfringens*. *Pakistan Veterinary Journal* 31 (3): 175-184.
- LÜCKSTÄDT, C. (2013). Effects of dietary sodium diformate in broilers – a performance analysis. *Proceedings of the 19th European Symposium on Poultry Nutrition*: 191.
- LÜCKSTÄDT, C. AND THEOBALD, P. (2009). Effect of a formic acid-sodium formate premixture on *Salmonella*, *Campylobacter* and further gut microbiota in broilers. *Proceedings and Abstracts of the 17th European Symposium on Poultry Nutrition*: 246.
- SELLE, P.H., HUANG, K.H. AND MUIR, W.I. (2004). Effects of potassium diformate inclusion in broiler diets on growth performance and nutrient utilisation. *Proceedings Australian Poultry Science Symposium*: 55-58.
- VAN DER SLUIS, W. (2000). *Clostridial* enteritis is an often underestimated problem. *World Poultry* 16: 42-43.