

Mechanisms of Resistance and Alteration of Chemical Compositions of the Potential Cash Crop Halophyte *Leptochloa fusca* L. Kunth under Salinity Stress



Tropentag 2016

Sayed Eisa¹, Nasr El-Bordeny¹, Ahmed Abdel-Ati², Mohamed Eid¹, Emad El-Din Abd El-Samad³, Sayed Hussin¹, Safwat Ali¹, Mohamed lotfy¹, Abdalla Masoud², Abd-El-Rahman El-Naggar¹, Mohamed Ebrahim¹

¹Ain Shams University (ASU), Faculty of Agriculture, Egypt; ²Desert Research Center, Plant Production Dept., Egypt; ³National Research Centre (NRC), Vegetables Crop Res. Dept., Egypt

1. Objectives

Rehabilitation dry-saline soil by using high potential halophyte. Saving the precious fresh water resources. Allow farmers to make productive use of dry-saline soil.

2. Introduction

Cash halophyte crops can grow using land and water unsuitable for other conventional crops and provide food, fodder, fuel, medicines, landscaping. One of these plants is Kallar Grass (*Leptochloa fusca* L. Kunth). It is a fast growing, perennial herbaceous, and can be utilized as forage, bio-reclamation of saline soil and carbon sequestration.

3. Materials and methods

• Potential yield of *Leptochloa fusca* under salt stress was evaluated using pot experiment.

• Leaf osmotic potential (-MPa), Potassium and sodium concentration, Transpiration rate ($\text{mmol m}^{-2}\text{S}^{-1}$) and Stomatal resistance ($\text{m}^2\text{S mol}^{-1}$) of *Leptochloa fusca* grown under 0.0%, 100, 200, 300 mM NaCl were determined.

• Evaluate the chemical composition of *Leptochloa fusca* as fodder crop under non saline soil and 100 mM NaCl .

4. Results




Fig. 1. Growth and development of *Leptochloa fusca* (A) and responses of shoot fresh weight to different NaCl salinity levels (b). The crossover of the dotted and solid lines reflects the limits of salinity resistance




Fig. 2. Leaf osmotic potential (-Mpa) of *Leptochloa fusca* grown under 0.0%, 100, 200, 300 mM NaCl

5. Conclusion

- The limits of salinity resistance for *Leptochloa fusca* was at 150 mM NaCl.
- *Leptochloa fusca* has the ability to regulate its osmotic potential under salinity stress.
- Has salt tolerance mechanism to avoid ion toxicity by replacement of potassium by sodium
- *Leptochloa fusca* could be utilized to rehabilitate dry-saline soil and provides high potential fodder crop




Fig. 3. Potassium and sodium concentration in *Leptochloa fusca* shoot at different NaCl salinity levels

Table 1. Effect of different NaCl levels on transpiration rate and Stomatal resistance of *Leptochloa fusca*

NaCl concentration (mM)	Transpiration rate ($\text{mmol m}^{-2}\text{S}^{-1}$)	Stomatal resistance ($\text{m}^2\text{S mol}^{-1}$)
Control	0.646 ^a	54 ^d
100	0.304 ^b	90 ^c
200	0.265 ^c	103 ^b
300	0.220 ^d	115 ^a

Table 2: Chemical composition of the *Leptochloa fusca* shoot at non saline soil and 100 mM NaCl.

NaCl level (mM)	Pot experiment	
	Control	100
Dry matter	90.55	90.79
Constituents, (% on DM basis)		
Organic matter	90.58	88.75
Ash	9.42	11.25
Crude protein	5.16	6.48
Crude Fiber	38.94	32.58
Ether extract	3.36	4.68
Nitrogen free extract	43.12	45.02
Cell wall constituents, % on DM basis		
Neutral-detergent fiber	76.21	72.49
Acid-detergent fiber	62.49	60.91