

# Increasing cowpea productivity by combining rock phosphate and arbuscular mycorrhizal fungi inoculation in sub-Saharan Africa

Kanako Suzuki<sup>1</sup>, Christian Fatokun<sup>2</sup>, and Ousmane Boukar<sup>3</sup>  
International Institute of Tropical Agriculture, <sup>1</sup>Lusaka, Zambia; and <sup>2</sup>Ibadan and <sup>3</sup>Kano, Nigeria

## Introduction

- The P deficiency negatively affects the formation of nodule (1, 2) and the symbiotic fixation of N<sub>2</sub> gas by limiting growth and survival of rhizobia (1, 3) in legume crops.
- According to our previous study, P and N uptake by cowpea are significantly correlated ( $r = 0.845$ ,  $P < 0.01$ ) (4).
- Additionally, P uptake has significant and high positive correlation with shoot dry weight (SDW) at 8 weeks after planting (WAP). This is a good indicator of P uptake in cowpea (4).
- Optimum amount of rock P application on cowpea was identified as 57 mg P kg<sup>-1</sup> (4).
- In sub-Saharan Africa, rock P deposit exists widely (5); thus, it can be used as alternative P source of chemical fertilizer owing to increase of P fertilizer price.
- Arbuscular mycorrhizal fungi (AMF) are known to help promote P and water uptake by crops by elongating their hyphae (6).
- Our study focused on the use of rock P and AMF inoculation to enhance P uptake by cowpea.

## Materials and Methods

- ◆ **Pot test 1** was conducted to verify the effect of AMF inoculation on cowpea shoot dry weight with 0 and 30 mg P kg<sup>-1</sup> as KH<sub>2</sub>PO<sub>4</sub>.
- ◆ **Pot test 2** was conducted to verify the effect of co-application of rock P (60 mg P kg<sup>-1</sup>) and AMF inoculation.
- ◆ **Pot test 3** was conducted to identify the optimum amount of rock P application for AMF inoculation using 20, 40, and 60 mg P kg<sup>-1</sup>.
- ◆ **Pot test 4** was conducted to identify the effect of co-application of rock P (40 mg P kg<sup>-1</sup>) and AMF inoculation on cowpea drought tolerance.

### Soil

- Pot tests 1, 2, and 3 were conducted using low P (approx. 1.06 ppm) soil collected from Fashola village (N07°53'718", E003°45'773") in Nigeria.
- In pot test 4, low P and N soil that was made mixing sand and clay collected in IITA-Nigeria. The available P was 0.48 ppm and total N was 0.004%. The percentages of sand, silt, and clay were 84, 5, and 11%, respectively.


### Nutrients, AMF and irrigation

- Rock P from Togo was along with AMF strain, *Glomus intraradices*.
- In all pot tests, basal application of 50 K (KCl), 50 Mg (MgSO<sub>4</sub> 7H<sub>2</sub>O), 5 Zn (ZnSO<sub>4</sub> 7H<sub>2</sub>O), 10 Mn (MnCl<sub>2</sub> 4H<sub>2</sub>O), 5 Cu (CuSO<sub>4</sub> 5H<sub>2</sub>O), 5 Mo ((NH<sub>4</sub>)<sub>6</sub>MO<sub>7</sub>O<sub>24</sub> 4H<sub>2</sub>O)mg kg<sup>-1</sup> soil (7) was employed.
- De-ionized water was used for irrigation to keep the soil water content at approx. 50% of field capacity in pot tests 1, 2, and 3.
- In pot test 4 (normal irrigation treatment), soil water content was kept at approx. 50% of field capacity. Drought treatment was irrigated 50 ml once per week from 3 WAP to 7 WAP. Water content under drought condition was less than 25% of field capacity.

## Results and Discussion

### Test 1: Can AMF inoculation help increase SDW in cowpea?


- Infection rate of AMF under zero P application was approx. 30-50% in all cowpea genotypes and maize, but under 30 mg P kg<sup>-1</sup> was almost 0%.
- Focusing on zero P application, the SDWs of cowpea genotypes with AMF inoculation were higher than without AMF inoculation (Fig. 1).
- **AMF inoculation does not work if there is high available P in soil. But under low P condition, AMF can help to increase SDWs of cowpea.**



**Figure 1: Shoot dry weight (SDW) at 8 weeks after planting (WAP) without (-) or with (+) arbuscular mycorrhizal fungi (AMF) under the treatment of 0 (0P) or 30 mg P kg<sup>-1</sup> (30P) as KH<sub>2</sub>PO<sub>4</sub> in 6 cowpea genotypes and a maize variety**  
Letters denote the significant differences at  $P < 0.05$  by Tukey's method compared to no-inoculated plant in each genotype under each P level. The vertical bar is the standard error. The yellow color represents the value of maize.

### Test 2: Can co-application of rock P and AMF application increase SDW of cowpea ?


- Cowpea genotype, Sanzi, only showed significantly ( $P < 0.05$ ) higher SDW under co-application of rock P and AMF inoculation than under rock P only (Fig. 2).
- The SDWs of other 14 cowpea genotypes (except Sanzi) under co-application of rock P and AMF inoculation were slightly higher than under rock P only (Fig. 2).
- **These results indicated that rock P application at 60 mg P kg<sup>-1</sup> is too high for AMF inoculation to work.**



**Figure 2: Shoot dry weight (SDW) at 8 weeks after planting (WAP) without (-) or with (+) arbuscular mycorrhizal fungi (AMF) inoculation under the treatment of 0 and 60 mg P kg<sup>-1</sup> as RP in 15 cowpea genotypes**  
The vertical bar represents the standard error. Letters denote the significant differences at  $P < 0.05$  by Tukey's method compared to 0 mg P kg<sup>-1</sup> without AMF inoculation.

### Test 3: How much of rock P is optimum for AMF inoculation to work?


- Mean of SDWs from the data of 6 cowpea genotypes was significantly higher under the treatment of 20 mg P kg<sup>-1</sup> compared to zero application.
- Significant differences among 20, 40, and 60 mg P kg<sup>-1</sup> were not identified.
- **The result explains that 20 mg P kg<sup>-1</sup> was optimum amount of rock P application for cowpea with AMF inoculation.**



**Figure 3: Means of SDWs at 8 weeks after planting (WAP) under the different RP application amounts; 0, 20, 40, and 60 mg kg<sup>-1</sup> with arbuscular mycorrhizal fungi (AMF) inoculation in 6 cowpea genotypes**  
The vertical bar represents the standard error. Letters denote the significant differences at  $P < 0.05$  by Tukey's method compared to 0 mg P kg<sup>-1</sup>.

### Test 4: Can co-application of rock P and AMF inoculation increase drought tolerance?

- TVu-7778 is a susceptible genotype to drought stress (8). However, young plant survived with co-application of rock P and AMF inoculation.
- Flowering of survived plants started from co-application of rock P and AMF inoculation and sole AMF inoculation at 10 WAP.
- **Co-application of rock P and AMF inoculation has high possibility of inducing drought tolerance of cowpea.**



**Photo 1: Responses of cowpea genotypes, TVu-7778 to drought stress under the different treatments (7WAP)**  
OP: zero P application, 40P: rock P application as 40 mg P kg<sup>-1</sup>, +AMF: with AMF inoculation, -AMF: without AMF inoculation.

## Conclusion

- Inoculation of AMF such as *Glomus intraradices* can increase SDW at 8 WAP of cowpea under low P condition. Therefore, it can contribute to increase P uptake by young cowpea plant.
- For AMF inoculation to work, we need to apply at least 20 mg P kg<sup>-1</sup> as rock P.
- Co-application of rock P and AMF inoculation has high possibility of inducing drought tolerance in cowpea.

## Acknowledgements

The authors gratefully appreciate the support provided by the Ministry of Foreign Affairs (MOFA) of Japan in the conduct of this study. We would also like to thank the staff of the Cowpea Breeding Unit at IITA for their invaluable help in this research work.

## References

- Drevon, J.J., Hartwig, U.A. 1997. Phosphorus deficiency increase the argon-induced decline of nodule nitrogenase activity in soy bean and alfalfa. *Planta* 201, 463-469.
- Kouas, S., Labidi, N., Debez, A., Abdell, C. 2005. Effect of P on nodule formation and N fixation in bean. *Agron. Sustain. Dev.* 25, 389-393.
- O'Hara, G.W., Boonkered, N., Dilworth, M.J. 1988. Mineral constraints to nitrogen fixation. *Plant Soil* 108, 93-110.
- Suzuki, K., Fatokun, C., Boukar, O. 2016. Abstract in book of abstracts pp 162-163, Pan-African Grain Legume & World Cowpea Conference, February 28 – March 4, Livingstone, Zambia.
- Nakamura, S., Fukuda, M., Nagumo, F., Tobita, S. 2013. Potential utilization of local phosphate rocks to enhance rice production in sub-Saharan Africa. *JARQ* 47, 353-363.
- Udaiyan, K., Gowsalya, Devi A.P., Chitra, A., Greep, S. 1997. Possible role of arbuscular mycorrhizal (AM) fungi on drought tolerance in *Vigna unguiculata* subsp. *unguiculata* (L.) Walp and *Leucaena latifolia* L., Pertanika. *J Trop. Agric. Sci.*, 20, 135-146.
- Kolawole, G.O., Tian, G., Singh, B.B. 2002. 4.7 Differential response of cowpea lines to application of P fertilizer. Challenges and opportunities for enhancing sustainable cowpea production (eds.) Fatokun, C.A., Tarawali, S.A., Singh, B.B., Kormawa, P.M., Tamo, M., International Institute of Tropical Agriculture, Ibadan, Nigeria. pp 319-328.
- Singh B.B. and Matsui T. 2002. 4.5 Cowpea varieties for drought tolerance. pp 287-300 in reference 7.