

# Agronomic evaluation of a collection of *Cenchrus ciliaris* under tropical dry forest conditions

Gutiérrez Solís, Jhon Freddy<sup>1</sup>; Vivas Quila, Nelson José<sup>2</sup>; Peters, Michael<sup>1</sup>; Morales, Sandra<sup>2</sup>; Hincapié Carvajal, Belisario<sup>1</sup>

<sup>1</sup>International Center for Tropical Agriculture (CIAT), Tropical Forages Program. <sup>2</sup>University of Cauca, Agricultural Sciences Department, Colombia.

CONTACT: [j.f.gutierrez@cgiar.org](mailto:j.f.gutierrez@cgiar.org)

## Introduction

Cattle farming is one of most important agricultural activities in Colombia and the main source of income for many farmers. In order to make the cattle sector more productive and resilient, there is a need to explore novel forage options that respond to adverse and changing climates in the tropics of Latin America and the Caribbean, in particular in warm dry environments.



Photo 1: Aerial photography of the trial


## Objective

Evaluate the agronomic performance of a collection of *Cenchrus ciliaris* under dry tropical forest conditions.

## Methodology



Altitude  
**500-800** masl



Average temperature  
**30-32° C**



Rainfall  
**1,200** mm/year

Once the collection of 20 accessions was established, an agronomic evaluation was carried out using the methodology of the International Network for the Evaluation of Tropical Pastures (RIEPT), which takes into account variables such as vigor, coverage, height, incidence of pests and diseases, flowering, green forage and dry matter production.



Photo 2: Plot of *Cenchrus ciliaris* (left). Aerial view of the plot (right).



Photo 3: Randomized complete block design, with four replications, and using for control commercial materials previously evaluated in the region (e.g. *Brachiaria brizantha* cv. Toledo).

## Results

The evaluations were carried out in two years during periods of maximum and minimum rainfall, which allowed to observe the performance year round resulting in the pre-selection of the following materials based on their dry matter production. The selected materials are complementary to the best control *Brachiaria brizantha* cv. Toledo which had a yields of 43.3 t ha<sup>-1</sup>year<sup>-1</sup>; *B. hybrid* cv. Cayman 40.31, and *Panicum maximum* cv. Mombasa 31.98, respectively. The genotypes evaluated with best yields were the accessions 6642, 15687 and 13299, with an average production of 39.59, 37.33 and 36.50 t ha<sup>-1</sup>year<sup>-1</sup>, respectively.




Figure 1: Performance of the materials taking into account the variables evaluated under the RIEPT methodology.

## Conclusions

The novel forage options selected are an excellent alternative for the dry tropical forest conditions and will allow diversification of options available to farmers, contributing to sustainable intensification and increased resilience.

## Acknowledgements

This work was done as part of the Program of development and use of forage resources in sustainable systems of bovine production for the department of Cauca - University of Cauca - Research Group NUTRIFACA. The work was funded by the Sistema General de Regalías (SGR), Colombia. We acknowledge the support of the Cauca Regional Government.

This work was undertaken as part of the CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS), which is carried out with support from CGIAR Fund Donors and through bilateral funding agreements. For details please visit [ccaafs.cgiar.org/donors](http://ccaafs.cgiar.org/donors). This work was undertaken as part of the CGIAR Research Program on Livestock. We thank all donors that globally support our work through their contributions to the CGIAR system



Poster presented at:  
**TROPENTAG 2018**

