



Tropentag, September 18-20, 2019, Kassel

“Filling gaps and removing traps
for sustainable resource management”

Land Use Effects Sediment Dynamics in the Headwaters of a Tropical Montane Forest in Western Kenya

JAQUELINE STENFERT KROESE¹, SUZANNE JACOBS², LUTZ BREUER³, JOHN QUINTON¹, MARIANA RUFINO¹

¹*Lancaster University, Lancaster Environment Center, United Kingdom*

²*Justus-liebig University Giessen, Centre for Intern. Development and Environmental Research (ZEU), Germany*

³*Justus-Liebig University Giessen, Inst. for Landscape Ecology and Resources Management (ILR), Germany*

Abstract

Land use effects sediment yield in the headwaters of the Sondu River basin in Kenya's largest remaining tropical montane forest, the Mau Forest Complex. Streams containing sediments originating from eroding cultivated land or unpaved roads can lead to on- and off-site effects that have economic and social impacts. The Sondu River drains into Lake Victoria, the second largest fresh water lake in the world, which is an important water resource for five countries and the source of the River Nile. East African streams are often characterised by high concentrations of suspended sediment, however, the availability of sediment yield data is severely limited. Within Kenya, there are only a few studies. Here, we report on a unique, four-year, high temporal resolution turbidity monitoring suspended sediment dynamics data set (10.2014–12.2018) which we use to assess temporal and spatial variability within three catchments (27–35 km²) under distinct land uses (natural forest, smallholder agriculture and tea/ tree plantations). As a surrogate for total suspended sediments, turbidity was calibrated by using an ex-situ suspended sediment increment suspension method. The outlet of the natural forest catchment had the lowest suspended sediment concentrations (34.5±76.4 mg L⁻¹) followed by the tea/ tree plantation (46.9±88.6 mg L⁻¹) and highest concentrations were observed at the outlet of the smallholder agricultural catchment (121.0±180.6 mg L⁻¹). Annual rainfall varied from 1554, 1730 to 1842 mm yr⁻¹ among the three catchments of the tea/ tree plantations, smallholder agriculture and natural forest, respectively. Sediment catchment yields showed seasonal variations in all three catchments with highest seasonal yields (4.2 – 24.1 t km⁻² month⁻¹) during the long rains (Apr - Jul) and lowest yields (0.4 - 0.5 t km⁻² month⁻¹) during the dry season (Jan - Feb), closely followed by discharge and rainfall patterns during the four year observation period. The high-resolution data set showed that the catchment under smallholder agriculture had the highest sediment yields when compared to the catchments under tea/ tree plantation and natural forest (106.0, 40.3 and 20.4 t km⁻² year⁻¹, respectively). The study indicated the importance of keeping natural forest ecosystems and managing agricultural systems to protect water resources from sediment pollution.

Keywords: Land use, suspended sediment yield, temporal variation, tropical mountain catchments, spatial variation