

Agronomic evaluation of 15 *Centrosema* sp. accessions in time of maximum and minimum rainfall in Colombia

Sotelo M; Peters M; Arango J; Cardoso JA.
Alliance Bioversity-CIAT, Tropical Forages Program
CONTACT: j.a.cardoso@cgiar.org

Introduction

- Worldwide, livestock is one of the most important economic activities for the economy of developing countries.
- In fact, it is the crop with the largest land use on the planet (around 3,400 million hectares for livestock feed).
- Much of the livestock sector is managed in grazing systems.
- Often native or naturalized pastures generate seasonal food shortages at critical times.

Objective

Agronomic evaluation of 15 accessions of *Centrosema* sp. obtained from the Genetic Resources Bank of The Alliance Bioversity International and CIAT, in Palmira, Colombia.

Methodology

- Data was collected for 14 months, between 2020 and 2021, including periods of maximum and minimum precipitation.
- The size of the plots was 9m² with three repetitions per treatment.
- Cuttings were carried out at different regrowth ages:
 - First cutting: 35 days
 - Second cutting: 42 days
 - Third cutting: 49 days after cutting.
- Evaluated variables: Agronomic parameters (coverage, vigor, height; Data on these parameters are not presented) and Dry matter production (DM).

Results and analysis




Figure 1. Biomass production under different recovery ages. Source: own elaboration.

Forage production

- The best accession when averaging biomass production based on DM in the two climatic seasons was CIAT 15474, surpassing control CIAT 15160 by 0.57 ton DM ha⁻¹.
- The highest forage production based on DM occurred in the second cutting with 7.12 ton DM ha⁻¹ for the CIAT 15474 accession, averaging the two climatic seasons.
- Centrosema* sp. It is a suitable material to be used as a forage alternative for animal feed according to biomass production under the soil and environmental conditions of Valle del Cauca in Colombia.
- Although the best age for harvest was the second cut, it is necessary to check the information obtained with an analysis of the nutritional quality of the materials.

Conclusions

- » Diversification in the inclusion of forage materials for animal feed, through the development of forage legumes, will provide multiple benefits at a productive and environmental level, improving the availability of food in critical times, positively impacting parameters such as weight gain of animals.
- » The Forage Network project generated critical information to identify promising materials that can be used in future experiments to mitigate greenhouse gas emissions in the tropics.

Further reading

Heinritz SN; Hoedtke S; Martens S; Peters M; Zeyner, A. 2012. Evaluation of ten tropical legume forages for their potential as pig feed supplement. Livestock Research for Rural Development 24, #7. <http://www.lrrd.org/lrrd24/7/hein24007.htm>
Peters M; Franco LH; Schmidt A; Hincapié B. 2011. Especies forrajeras multipropósito: opciones para productores del trópico Americano. Publicación CIAT no. 374. Centro Internacional de Agricultura Tropical (CIAT); Cali, CO. <https://hdl.handle.net/10568/54681>

PRINTED WITH SOLVENT INKS This poster is licensed for use under the Creative Commons Attribution 4.0 International license (CC BY 4.0) 2022-08. Design: A Yedra/CIAT. Photo: M Sotelo/CIAT.

Acknowledgements

This work was undertaken as part of the AGROSAVIA Forages Network and the CGIAR Initiatives on Livestock, Climate and System Resilience (LCSR); Sustainable Animal Productivity for Livelihoods, Nutrition and Gender Inclusion (SAPLING). We thank all donors that globally support through their contributions to the CGIAR system.

