




# Benefits of *Leucaena diversifolia* in grazing steer's diet: performance, methane and fatty acids

Isabel C. Molina-Botero<sup>1</sup>; Daniel Mauricio Villegas<sup>1</sup>; Alejandro Montoya<sup>1</sup>; Johanna Mazabel<sup>1</sup>; Mike Bastidas<sup>1</sup>; Alejandro Ruden<sup>1,2</sup>; Hernan Gaviria<sup>3</sup>; Juan David Peláez<sup>3</sup>; Julián Chará<sup>4</sup>; Enrique Murgueitio<sup>4</sup>; Jon Moorby<sup>5</sup>; **Jacobo Arango<sup>1</sup>**

<sup>1</sup>International Center for Tropical Agriculture, Tropical Forages Program, Colombia; <sup>2</sup>University of Florida, USA; <sup>3</sup>Asobrangus Comercial ANGUSAZUL<sup>®</sup>, Colombia; <sup>4</sup>Center for Research on Sustainable Agricultural Production Systems - CIPAV, Colombia; <sup>5</sup>Aberystwyth University, UK.

Contact: j.arango@cgiar.org



## Introduction

Livestock production contributes significantly to global greenhouse gas emissions (UNEP and ACC, 2021). Some strategies to reduce enteric methane (CH<sub>4</sub>) emissions propose modulating ruminal fermentation by providing legumes rich in secondary compounds.

*Leucaena diversifolia* is a legume species that has received little attention in terms of its nutritive value, CH<sub>4</sub> emissions, and impact on meat quality. To address this gap, a study was conducted to compare the performance, CH<sub>4</sub> emissions, and fatty acid content of steers grazing on a monoculture of tropical grass *Urochloa* hybrid cv. Cayman versus a combination of Cayman and *L. diversifolia*.



## Methodology

**Location:** The experiment was conducted at the International Center for Tropical Agriculture (CIAT) located in Valle del Cauca, Colombia.

**Animals and diets:** Over a period of 15 months, 14 Angus crossbred steers were used in the study, with half of them grazing only Cayman grass (**CG**, *Urochloa* hybrid cv. Cayman-CIAT BR02/1752) and the other half grazing on a combination of Cayman and *L. diversifolia* ILRI 15551 (**SPS**) at a ratio of 74:26.

Table 1. Mean (+/- SD) nutritional composition of the diets offered to the steers during the experiment.

Variable	DM g/kg	CP g/kg DM	IVDMD g/kg DM	Total phenols g/kg DM	Tannins g/kg DM
CG	242±1.6	64±1.4	467±2.5	3.30	0.10
SPS	232±3.4	135±0.6	639±2.9	17.42	12.40

**SPS:** 74% *Urochloa brizantha* cv. Toledo and 26% *Leucaena diversifolia*; **CG:** Cayman grass; **DM:** dry matter; **CP:** crude protein; **IVDMD:** *in vitro* DM digestibility.

**Evaluations:** The animals were weighed every 4 months throughout the experiment. At the end of the trial, enteric CH<sub>4</sub> emissions were measured for one month using the polytunnel technique (Murray et al., 2001). Subsequently, the animals were slaughtered to extract a subsample of the *longuissimus dorsi* muscle to quantify fatty acid esters (Mojica-Rodríguez et al., 2017).



## Results

Animals grazing Cayman grass plus *L. diversifolia* consumed 1.22 times more dry matter than animals ingesting grass alone ( $P \leq 0.05$ ). This difference was doubled when calculating daily crude protein intake ( $P \leq 0.05$ ). At the end of the experiment, animals consuming the **SPS** diet weighed 63 kg more than animals consuming **CG** ( $P \leq 0.05$ ). On the other hand, steers consuming *L. diversifolia* emitted 15% less net enteric CH<sub>4</sub> than animals on **CG** ( $P \leq 0.05$ ). Likewise, in our study the omega 6: omega 3 ratio was lower in animals consuming this legume ( $P \leq 0.05$ ).

Table 2. Forage intake, animal productivity, methane emissions and meat quality of steers fed with and without *L. diversifolia*

Variable	Treatment		SEM	P-value
	SPS	CG		
<b>Intake</b>				
DM (kg/d)	8.571	6.995	0.514	0.0082
CP (g/d)	847.0	396.9	27.63	0.0001
<b>Animal productivity and emissions</b>				
Final LW (kg)	535.3	472.3	10.22	0.0410
Net methane emissions (g/d)	142.4	167.6	5.425	0.0069
<b>Fatty acids</b>				
Omega-6: omega-3 ratio	2.59	2.91	0.223	0.0173

**SPS:** 74% *U. brizantha* cv. Toledo or *U. hybrid* cv. Cayman and 26% *L. diversifolia*; **CG:** 100% *U. hybrid* cv. Cayman or *U. brizantha* cv. Toledo; **SEM:** standard error of mean. **DM:** dry matter; **CP:** crude protein; **LW:** live weight.



## Conclusions

- The results of this study suggest that the inclusion of *L. diversifolia* improves nutrient intake, increases animal productivity, reduces daily enteric methane emission, and omega-6: omega-3 ratio in meat is reduced.

## References

- UNEP and CAC, 2021. Global methane assessment: Benefits and costs of mitigating methane emissions. United Nations Environment Programme and Clean Air Coalition, Nairobi. [Cited 2023 May]. Available from: <https://www.unep.org/resources/report/global-methane-assessment-benefits-and-costs-mitigating-methane-emissions>
- Murray, P.J., Gill, E., Balsdon, S.L., Jarvis, S.C., 2001. A comparison of methane emissions from sheep grazing pastures with differing management intensities. *Nutr. Cycl. Agroecosyst.* 60, 93-97
- Mojica-Rodríguez, J. E., Castro-Rincón, E., Carulla-Fornaguera, J., Lascano-Aguilar, C. E., 2017. Efecto de la edad de rebrote sobre el perfil de ácidos grasos en gramíneas tropicales. *Ciencia y Tecnología Agropecuaria*, 18(2), 217-232

## Acknowledgments

We thank all donors that globally support the work of the OneCGIAR initiative on Livestock and Climate. Advancing sustainable forage-based livestock production systems in Colombia (CoForLife). We also thank Angus Azul<sup>®</sup> for the acquisition of animals and CIPAV for the support received during the execution of the experiment.



Poster prepared for:  
**Tropentag 2023**  
September 20-22, 2023  
Berlin (Germany)