

## **GROWTH AND ECONOMIC PERFORMANCE OF BROILER CHICKENS FED TURMERIC RHIZOME POWDER (TRP) (*CURCUMA LONGA* LINN)**

Enyenihi, G. E., Ubak E, Ekpo; Dan, D. A., and Ogbuzuru, E. N.

Corresponding author's email: [gloryeenyenihi@uniuyo.edu.ng](mailto:gloryeenyenihi@uniuyo.edu.ng),

Corresponding author's Phone: (+234) 8061513068

Department of Animal Science, University of Uyo, Akwa Ibom State, Nigeria.

### **Abstract**

This study was conducted at the University of Uyo Teaching and Research Farm to evaluate the impact of turmeric rhizome powder (TRP) supplementation on growth performance of broiler chickens and cost efficiency of broiler production. A total of 120 Ross 308 strains of broiler chickens were divided into four treatment groups, which were further replicated three times with 10 birds each per replicate in a Completely Randomized Experimental Design. The experimental groups were made up of the birds on the control diet (T1), and three experimental groups that were fed diets with TRP supplementation at 200g (T2), 400g (T3), and 600g (T4) per 100kg of feed. The experiment lasted 8 weeks, during which data on weight gain, feed efficiency and cost efficiency were collected. Analysis of Variance (ANOVA) was adopted to detect significant differences between the treatment means while post hoc analysis using Duncan Multiple Range Test further differentiated the means. The results revealed that the birds on T3 diets (400g TRP) exhibited the highest weight gain (1508.33g), demonstrating the potential of TRP supplementation at that level of inclusion to enhance broiler growth performance. Furthermore, the cost analysis reveals that the cost per kilogram weight gain was least in the treatment fed 400g of TRP (₦959.00). Equally, the revenue and the gross margin was also highest in the treatment fed 400g of TRP (₦1885.00, ₦439.00). In contrast, exceeding this level of 400g supplementation, as observed in the T4 group (600g TRP), led to an increase in production costs without any proportional improvement in weight gain.

**Keywords:** broilers, additives, turmeric, curcuma and broiler economics.

### **Introduction**

High population growth and urbanization is driving the demand for meat and other high protein foods of animal origin in Nigeria and the rest of the developing world. As a means of achieving sufficient protein intake, broiler meat remains one of the chief sources that Nigerians rely on because of its availability as well as its popularity as a component of the menu of the generality of the populace (Anyanwu *et al.*, 2015). The poultry industry value chain in Nigeria had experienced a boom in recent time but currently challenged by several factors including cost of production (Mondal *et al.*, 2015). Average feed prices have at least doubled within the last one-year and a half, which has not only put many farmers out of business but has also reduced profit margins considerably for the resilient ones (Ettah *et al.*, 2021). High feed costs are driven by many factors; notable amongst them is the reduced supply of major conventional feed ingredients such as maize and soya bean due to insecurity and insurgency ravaging the major grain producing areas of northeast and north central Nigeria (Anyanwu *et al.*, 2015).

In these areas, farmers are unable to access their farms, as a result, there appears to be more industrial demand for the little quantity of grains and pulses available (Ettah *et al.*, 2021). Consequently, the high cost of feed has necessitate the need for research for alternative feed ingredients and a means of improving the feed conversion ratio. This has led to the emergence of numerous unconventional feedstuff and increased demand for organic feed additives in the market,

in the form of probiotics, antibiotics and growth promoters. Growth promoters are chemical and biological substances, which are added to livestock feed to improve feed utilization and growth for better production and financial results (Mondal *et al.*, 2015). Their mode of action could be through better appetite, improved feed conversion, stimulation of the immune system and increased vitality (Malik *et al.*, 2022).

Under intensive chicken production systems, feed additives and or growth promoters are often used to suppress or eliminate harmful microorganisms in the intestines and to improve growth and performance. However, synthetic growth promoters and hormones have disadvantages of high cost, adverse side effect on health of birds and residues in meat and development of antibiotic resistance in microbes (Mondal *et al.*, 2015). In view of the recent increase in the demand for organically produced food free from antibiotics, research on the use of natural plant ingredients to replace the synthetic products has received more attention. Alternative substances and strategies for animal growth promotion and disease prevention have been and are being investigated. Some phytogenic and herbal products which have received high acceptability among consumers in recent times have been targeted (Nouzarian *et al.*, 2011).

Amongst these lists of natural supplements capable of eliciting physiological responses when added to the broiler diets, include members of the Zingiberaceae family such as ginger and turmeric. Other researchers have reported on other additives such as *Moringa oleifera* (Anyanwu *et al.*, 2020), garlic, scent leaf and antibiotic (Odoemelam *et al.*, 2017). Turmeric (*Curcuma longa*) is a food spice used mainly for its human nutritional and medicinal potentials. It is a rhizomatous herbaceous perennial plant of the Zingiberaceae family and has its native origin in southeastern India where it is known as golden spice of India and has a very long history of medicinal use, dating back nearly 4000 years (Akaberi *et al.*, 2021).

It is used not only as a principal spice but also as a component in religious ceremonies (Samarasinghe *et al.*, 2003). Its brilliant yellow colour gives it the name. "Indian saffron." (Prasad and Aggrawal, 2011). According to Prasad and Aggrawal (2011), India is the largest producer of turmeric and supplies 94% of the world's demand. The main active substance of turmeric extract is identified as curcumin, a strong antioxidant and a peptide named turmerin (Samarasinghe *et al.*, 2003). Turmeric has been shown to have several biological effects in humans, exhibiting anti-inflammatory (Holt *et al.*, 2005), antioxidant (Iqbal *et al.*, 2003) and hypolipidaemic (Ramirez-Tortosa *et al.*, 1999) activities. It has also been suggested that turmeric possess hepato-protective, anti-tumor, antiviral and anticancer activities (Polasa *et al.*, 1991). Reports also exist indicating that it has been used for the treatment of gastrointestinal and respiratory disorders in humans (Anwarul *et al.*, 2005). However, its effect on broiler chickens at very high levels of inclusion has not been widely reported. It is in the light of these physiological effects and economic importance of turmeric (*Curcuma longa*), that this study was conducted to evaluate the performance of raw turmeric powered (*Curcuma longa*) rhizome supplemented diet meal on broiler chickens.

## **Materials and Method**

### **Experimental site**

The experiment was carried out at the Poultry unit of the Department of Animal Science Teaching and research Farm, University of Uyo. The area falls within the tropical rainforest zone of Nigeria, with two distinct seasons, rainy season (March – Mid November) and dry season (November - March) it has an average rainfall of 2200mm to 3500mm. The temperature of the area ranges from 26 to 28°C (University of Uyo Meteorological station, 2018).

### Procurement and processing of Experimental Materials

The turmeric rhizomes that were used for this experiment were purchased from a vendor in a local market in Uyo metropolis. The rhizomes were washed, peeled and slice into smaller sizes after which the sliced chops were washed and sun dried. When they were dried enough, the rhizomes were ground using mechanical blenders into fine powders. The powders were further stored using tightly sealed containers.

### Experimental Diet

Four (4) straight experimental broiler diets were formulated for the four treatments for the whole period of the experiment. The diets were formulated with the supplementation of Turmeric rhizome Powder (TRP) that was locally prepared at different levels. Treatment 1 representing the control had no TRP while T2, T3, and T4 contained 200g, 400g and 600g per 100Kg of the formulated feed. The other nutritional ingredients were adjusted to meet the nutrient requirement of the broilers at all the ages of the experiment. The comprehensive ingredient composition and the calculated chemical composition of the formulated diets are presented in Table 1.

**Table 1: Composition of experimental diets (straight diet) using varying levels of turmeric rhizome powder (TRP).**

<b>Ingredients</b>	<b>T<sub>1</sub> (0g)</b>	<b>T<sub>2</sub> (200g)</b>	<b>T<sub>3</sub> (400g)</b>	<b>T<sub>4</sub> (600g)</b>
Maize	52.00	52.00	52.00	52.00
Full fat soya	20.00	20.00	20.00	20.00
Ground nut cake	11.00	11.00	11.00	11.00
Fish meal	3.00	3.00	3.00	3.00
Wheat offals	5.20	5.20	5.20	6.20
Palm kernel meal	5.00	5.00	5.00	5.00
Bonemeal	3.00	3.00	3.00	3.00
Common Salt	0.25	0.25	0.25	0.25
Vitamin /mineral premix	0.25	0.25	0.25	0.25
Lysine	0.20	0.20	0.20	0.20
Methionine	0.10	0.10	0.10	0.10
TRP	0.00	0.02	0.04	0.06
<b>Total</b>	<b>100.00</b>	<b>100.02</b>	<b>100.04</b>	<b>100.06</b>
<b>Calculated Analysis</b>				
Crude protein (%)	22.13	22.13	22.13	22.13
Metabolizable Energy ME (kcal/kg)	2985.2	2985.22	2985.22	2985.22
Crude fat (%)	26.85	6.85	6.85	6.85
Crude fibre (%)	3.76	3.76	3.76	3.76
Calcium (%)	1.10	1.10	1.10	1.10
Phosphorus (%)	0.45	0.45	0.45	0.45

Vitamin premix = vitamin A, D3, E,K, B1, B2,B6, B12, Nacin, Patothemic acid, Folic acid, Biotin, Choline Chloride, Manganese, Zinc, iron, Copper, iodine, selenium, cobalt and Antioxidants.

## Experimental birds and Design

A total of 120 unsexed Ross 308 broiler strains were used for the experiment. The experimental birds were purchased from a vendor in Uyo Metropolis. The birds were managed in a deep litter system with wood shavings as the litter material during the brooding and rearing phases. The experiment lasted for 56 days. Water and feed were provided for the experimental birds *ad libitum*. The birds were fed daily during the early hours of the day and leftovers were weighed every day to ascertain the daily feed intake of each replicate. Routine vaccination and drugs were administered to the birds in accordance with the prescriptions of the veterinary doctor.

The chicks were randomly divided into four (4) treatment groups (T1, T2, T3, and T4), with thirty (30) birds each per treatment. Each of the four treatments was further divided into three (3) replicates with ten (10) birds per replicate and the birds were randomly assigned to each of the formulated diets in a Completely Randomized Design (CRD).

## Data Collection

The data collected for these experiments were initial weight, total feed intake, daily feed intake, weekly weight gain, final weight and FCR. The process of obtaining these data is explained in the below:

The experimental birds were fed *ad libitum* on daily basis. Daily feed intake was calculated by subtracting the daily leftover from the quantity of feed given the previous day. The total feed intake therefore was calculated by summing all the daily feed intake from the first to the last day of the experiment.

On the first day of the experiment, the initial weight of the experimental birds was taken by weighing them individually according to their replicates. This same process was carried out weekly to obtain their weekly weight and weekly weight gain. On the last day of the experiment, the final weight of the experimental birds was recorded, and total weight gain was obtained by subtracting the initial weight from the final weight.

Feed conversion ratio is the measure of the ratio of the total feed intake that was used by the chicken to gain weight. It was calculated by dividing the total feed intake by the total weight gained by the chicken. Where:

$$\text{FCR} = \frac{\text{Total feed intake}}{\text{Total weight gain}}$$

The data for analysis of cost for this experiment was obtained using standard economic principles. The following data were collected: cost of feed consumed, cost/kg weight gain, total cost of production, revenue, and gross margin.

The cost of feed consumed per chicken was calculated by calculating the total cost of the total feed intake per chicken for all the replicates and treatments.

The cost per kilogram weight was obtained by dividing the cost of production per chicken by the total weight gain in kilogram.

$$\text{Cost per Kilogram Weight Gain} = \frac{\text{Total Production Cost}}{\text{Weight Gain}}$$

Revenue was calculated by multiplying the price per kilogram of chicken by the quantity of chicken produced.

The gross margin was calculated by subtracting the cost of producing the broiler chicken from the revenue generated from the sales of the broiler chicken.

## Data Analysis

Data generated from the experiment were subjected to analysis of variance (ANOVA) according to Steel and Torie (1980). ANOVA detected the treatment effects while means were separated using the Duncan Multiple Range Test.

## Result and Discussion

### Effect of dietary supplementation of TRP on the growth performance of broiler chickens.

The effect of supplementing broiler diets with turmeric rhizome powder (TRP) on their growth performance is presented in Table 2. The table shows that TRP supplementation on broiler diets had a significant effect ( $P < 0.05$ ) on the growth parameters measured beginning from initial weight up to the FCR except feed intake. The findings showed that weight gain for the chickens in the T3 (400g) group recorded the highest weight gain (1508.33g), followed by the T2 (200g) group (1390.92g), the T1 (control) group (1375.00g). However, those on T4 (600g) group recorded the lowest weight gain (1225.00g). While the highest weight was recorded in T3, it was statistically different ( $p < 0.05$ ) from the T4 but was statistically similar ( $p > 0.05$ ) with those of the T1 and T2.

Equally, the same trend was observed in the daily weight gain of the broiler chickens where the T3 (400g) group was also the highest (35.91g), followed by the T2 (200g) group (33.12g), the T1 (control) group (32.74g), and then the T4 (600g) group (29.16g). Just like in total and daily weight gain, the final weight gain of the experimental birds was found to also be highest in T3 which is significantly different from T4 but was similar to T1 and T2.

The feed conversion ratio (FCR) is a measure of how much a bird converts the consumed feed into flesh and in this present study, the best FCR (1.92), was recorded in the broiler chickens in the T3 (400g) group followed by the T2 (200g) group (2.01), the T1 (control) group (2.18), and the T4 (600g) group (2.43). The difference that existed between the groups in the FCR also follows the trend which is reported in the daily, total weight and final weight. The T3 was highest and statistically different ( $p < 0.05$ ) from T4 but was statistically similar ( $p > 0.05$ ) to T1 and T2.

This findings from this study suggests that supplementing broiler chicken diets with turmeric rhizome powder at a level up to 400g/100kg can improve their growth performance and feed utilization. From these figures, it can be inferred that feeding TRP as a dietary supplement to broilers is a beneficial practice that can be adopted in practical farming and chicken production. However, the study also showed that feeding TRP beyond this threshold of 400g/100kg was not beneficial.

The positive outcome achieved in this study is similar to the outcomes observed by Samarasinghe *et al.* (2003); Riasi *et al.* (2012); Hussien (2013); Kafi *et al.* (2017); Guil-Guererro *et al.* (2017); and Oke (2018) except that some reported an optimal ratio of supplementation that are higher than 400g to 100kg of feed. In the study of Samarasinghe *et al.*, (2003), two bioassays were conducted to assess the effectiveness of turmeric root powder and mannan-oligosaccharides (MOS) as antibiotic alternatives for broilers, and it was found turmeric (1g/kg) in the first trial generally improved broiler weight gain by 5.3%, in the second trial, it also significantly increased weight gain by 15.1%, respectively. In another study where ginger and turmeric powders were fed to broiler chickens at the rate of 0.50% and 0.75% of the diet respectively, both were found to exert positive impact on the weight gain and FCR conversion Ratio, however, garlic was reported to have performed better than the turmeric Kafi *et al.* (2017).

Riasi *et al.* (2012), also found that turmeric rhizome powder improved the Feed Conversion Ratio of older laying hens. Supplementing broiler diets with 7 g/kg of Turmeric powder have also positively influenced growth performance, reduced abdominal fat, and lowered serum cholesterol and triglyceride levels in broiler chickens (Hussien, 2013). This study differs from the present study because 7g/kg is higher than the 4g/kg which we found optimal.

The positive effect of turmeric rhizome powder on the physiological responses of broiler chickens was also consistent in the study of Oke (2018), as the supplementation of 8g turmeric per kilogram of diet was found to improve the final weight, weight gain and feed conversion ratio, all of which are key indices of a broiler's performance. But once again this study differs from our findings as the amount which Oke (2018) reported to be optimal is double of ours and higher than 600/100kg which had a negative impact in our present study.

**Table 2: Growth Performance of broiler chickens fed diets supplemented with TRP**

Parameters	T1 (Control)	T2 (200g)	T3 (400g)	T4 (600g)	SEM
Initial Weight(g)	308.33 <sup>a</sup>	275.75 <sup>b</sup>	291.67 <sup>ab</sup>	273.00 <sup>b</sup>	5.60
Daily Feed intake (g)	71.68	65.88	69.02	70.89	2.80
Total Feed intake (g)	3010.63	2766.49	2898.88	2977.67	117.46
Total weight gain (g)	1375.00 <sup>ab</sup>	1390.92 <sup>ab</sup>	1508.33 <sup>a</sup>	1225.00 <sup>b</sup>	64.66
Daily weight gain (g)	32.74 <sup>ab</sup>	33.12 <sup>ab</sup>	35.91 <sup>a</sup>	29.16 <sup>b</sup>	1.54
Final Weight (g)	1683.33 <sup>ab</sup>	1666.67 <sup>ab</sup>	1800.00 <sup>a</sup>	1583.33 <sup>b</sup>	64.55
FCR	2.18 <sup>ab</sup>	2.01 <sup>ab</sup>	1.92 <sup>b</sup>	2.43 <sup>a</sup>	0.13

<sup>ab</sup> Means with different letters on the same row are significantly different from each other. FCR = Feed Conversion Ratio

### **Cost of producing broiler chickens fed diets supplemented with Levels of TRP**

The result for the analysis of the cost of production for this study is presented in Table 3. The result from this study, showed that supplementing broiler diets with graded levels of turmeric rhizome powder had a significant effect ( $P < 0.05$ ) on the cost of producing broiler chicken as well as on the revenue and gross margin generated from the production. The result showed that the control; T1 (0g) had the highest cost per kilogram weight gain (N1013.00), while T2 (200g) and T3 (400g) had the lowest cost per kilogram weight gain (N960.00 and N959.00, respectively) while T4 (600g) had a higher cost per kilogram weight gain than T2 and T3, but it was still lower than T1.

In the analysis for Gross margin, TRP had a significant effect ( $P < 0.05$ ), and it was found that T3 (400g) had the highest gross margin (N439.00), followed by T2 (200g) with N413.00, and T1 (0g) with N337.00. However, it was found that T4 (600g) had the lowest gross margin (N127.00). Additionally, the data for the revenue generated revealed a significant effect ( $P < 0.05$ ) of supplementing broiler diets with TRP. It was found that T3 (400g) had the highest revenue (N1885.00), followed by T2 (200g) with N1739.00, and T1 (0g) with N1719.00. Equally, just like in the gross margin, the T4 (600g) also generated the lowest revenue N1531.00).

From these figures, it can be inferred that feeding TRP as a dietary supplement to broilers is a beneficial practice that can be adopted in practical farming and chicken production. However, the study also showed that feeding TRP beyond a certain threshold may instill a negative economic impact. In this case, it was shown that 600g of TRP per 100kg of feeds does more economic harm than good to a farmer whose main aim is to maximize profit.

Scholarly, there is scarcity of literature that have studied the economic impact of turmeric rhizome powder supplementation on broiler production, but our study agreed to those of Kafi *et al.* (2017) who found that turmeric supplementation at 0.75% improved the economic viability of broiler chicken as well as their growth performance. Given that the growth performance and physiologic performance of broilers correlates positively with their profitability, this study agrees with some past literatures including Oke (2018) who found that TRP positively influences the physiological response and growth performance of broiler chickens. In a similar study conducted by Hussein *et al.* (2013), it was found that growth performance of broilers was higher in birds fed 7g TRP per kilogram feed.

**Table 3: Cost analysis of producing chickens fed diets supplemented with TRP**

Parameters	T1 (0g)	T2 (200g)	T3 (400g)	T4 (600g)	SEM
Cost per kg feed (₦)	459.11	479.11	499.11	519.11	60.11
Cost per kg weight gain (₦)	1013.00 <sup>ab</sup>	960.00 <sup>b</sup>	959.00 <sup>b</sup>	1146 <sup>a</sup>	50.40
Cost of feed consumed (₦)	1382.00	1325.00	1447.00	1404.00	35.50
Cost of Production (₦)	1382.00	1325.00	1447.00	1404.00	35.50
Gross Margin (₦)	337.00 <sup>ab</sup>	413.00 <sup>a</sup>	439.00 <sup>a</sup>	127.00 <sup>b</sup>	82.60
Revenue (₦)	1719.00 <sup>ab</sup>	1739.00 <sup>ab</sup>	1885.00 <sup>a</sup>	1531.00 <sup>b</sup>	80.80

<sup>ab</sup> Means with different letters on the same row are significantly different from each other. FCR = Feed Conversion Ratio

### Conclusion

Supplementing broiler diet with turmeric rhizome powder (TRP) improved their growth and feed efficiency, especially at a level of 400 grams per 100kg of feed. This amount of TRP also reduced production costs and increased profits, but higher levels had negative economic effects.

### References

- Akaberi, M., Sahebkar, A., and Emami, S. A. (2021). Turmeric and curcumin: from traditional to modern medicine. *Studies on Biomarkers and New Targets in Aging Research in Iran: Focus on Turmeric and Curcumin*, 15-39.
- Anuwarul, H. G., AbdulJabbar, S., Muhammed, N. G., and Kashif, M. (2005). Pharmacological basis for the use of turmeric in gastrointestinal and respiratory disorders. *Life Science*, 76(26), 3089-105
- Anyanwu, N. J., Ibe, C. E., Ekpe, I. I., and Etela, I. (2015). Effect of graded levels of inclusion of *Calapogonium muconoides* leaf meal on the performance of broiler finisher birds. *Tropical Animal Production Investigations*, 18(1), 60-67.

- Ettah, O. I., Igiri, J. A., and Ihejiamaizu, V. C. (2021). Profitability of broiler production in cross river state, Nigeria. *Global journal of agricultural sciences*, 20(1), 35-40.
- Guil-Guerrero, J. L., Ramos, L., Paredes, J. Z., Carlosama-Yépez, M., Moreno, C., and Ruales, P. (2017). Effects of turmeric rhizome powder and curcumin in poultry production. A review. *Journal of Animal and Feed Sciences*, 26(4), 293-302.
- Holt, P. R., Katz, S., and Kirshoff, R. (2005). Curcumin therapy in inflammatory bowel disease: a pilot study. *Digestive diseases and sciences*, 50, 2191-2193.
- Hussein, S. N. (2013). Effect of turmeric (*Curcuma longa*) powder on growth performance, carcass traits, meat quality, and serum biochemical parameters in broilers. *J. Adv. Biomed. Pathobiol. Res*, 3(2), 25-32.
- Iqbal, M., Sharma, S. D., Okazaki, Y., Fujisawa, M., and Okada, S. (2003). Dietary supplementation of curcumin enhances antioxidant and phase II metabolizing enzymes in ddY male mice: possible role in protection against chemical carcinogenesis and toxicity. *Pharmacology and toxicology*, 92(1), 33-38.
- Kafi, A., Uddin, M. N., Uddin, M. J., Khan, M. M. H., and Haque, E. (2017). Effect of dietary supplementation of turmeric (*Curcuma longa*), ginger (*Zingiber officinale*) and their combination as feed additives on feed intake, growth performance and economics of broiler.
- Malik, A. A., Busari, I. B., Idundun, S. O., Adebowale, Z. O., and Okoye, A. O. (2022). Effect of feeding *Jatropha tanjorensis* leaf meal as a growth promoter on the performance, carcass characteristics and sensory properties of broiler chicken. *Nigerian Journal of Animal Production*, 49(2), 180-190.
- Mondal, M. A., Yeasmin, T., Karim, R., Siddiqui, M. N., Raihanun-Nabi, S. M., Sayed, M. A., and Siddiky, M. N. A. (2015). Effect of dietary supplementation of turmeric (*Curcuma longa*) powder on the growth performance and carcass traits of broiler chicks. *SAARC Journal of Agriculture*, 13 (1), 188-199.
- Nouzarian, R., Tabeidian, S. A., Toghyani, M., Ghalamkari, G., and Toghyani, M. (2011). Effect of turmeric powder on performance, carcass traits, humoral immune responses, and serum metabolites in broiler chickens. *Journal of Animal and Feed Sciences*, 20(3), 389-400.
- Odoemelam, V. U., Unamba-Opara, I. C., Irondi, J. I., and Anyanwu, N. J. (2017). Performance and carcass quality of broilers fed dietary scent leaf, garlic, and antibiotic. *Nigerian Journal of Agriculture, Food and Environment*, 13(1), 188-193
- Oke, O. E. (2018). Evaluation of physiological response and performance by supplementation of *Curcuma longa* in broiler feed under hot humid tropical climate. *Tropical animal health and production*, 50, 1071-1077.
- Polasa, K., Sesikaran, B., Krishna, T. P., and Krishnaswamy, K. (1991). Turmeric (*Curcuma longa*)-induced reduction in urinary mutagens. *Food and Chemical Toxicology*, 29(10), 699-706.



- Prasad, S., and Aggarwal, B. B. (2011). Turmeric, the golden spice. *Herbal Medicine: Biomolecular and Clinical Aspects*. 2nd edition.
- Prasad, S., and Aggarwal, B. B. (2011). Turmeric, the golden spice. *Herbal Medicine: Biomolecular and Clinical Aspects*. 2nd edition.
- Ramirez-Tortosa, M. C., Urbano, G., López-Jurado, M., Nestares, T., Gomez, M. C., Mir, A., ... and Gil, A. (1999). Extra-virgin olive oil increases the resistance of LDL to oxidation more than refined olive oil in free-living men with peripheral vascular disease. *The Journal of nutrition*, 129(12), 2177-2183.
- Riasi, A., Kermanshahi, H., and Mahdavi, A. H. (2012). Production performance, egg quality and some serum metabolites of older commercial laying hens fed different levels of turmeric rhizome (*Curcuma longa*) powder. *Journal of Medicinal Plants Research*, 6(11), 2141-2145.
- Samarasinghe, K., Wenk, C., Silva, K. F. S. T., and Gunasekera, J. M. D. M. (2003). Turmeric (*Curcuma longa*) root powder and mannanoligosaccharides as alternatives to antibiotics in broiler chicken diets. *Asian-Australasian Journal of Animal Sciences*, 16(10), 1495-1500.
- Steel, R. G. D and J. H. Torrie (1980). *Principles and Procedures of Statistics*, McGraw-Hill Co. Inc. New York.
- University of Uyo Meteorological Station (2018).