

Towards a local system for screening child malnutrition

Cherif ISSIFOU, Bourama GUINDO, Waliou AMOUSSA HOUNKPATIN, Irene MITCHODIGNI, and Brigitte KAUFMANN

¹University of Abomey-Calavi, Faculty of Agricultural Sciences, Benin German Institute for Tropical and Subtropical Agriculture (DITSL), Witzenhausen, Germany

Introduction

- Screening is crucial in the fight against malnutrition
- But the conventional screening tools available are unfamiliar to rural caregivers.
- While caregivers are primarily responsible for child health and they have their own methods for detecting malnutrition in children.
- However, this local method and the way in which it is used are unknown to all comunity members and beyond the communities



Objective

To describe the local method used by caregivers to assess the nutritional status of their children



Study area

- Nikki et Banikoara/Benin
- Baatonu, Fulani and Gando ethnic groups

Approach: Co-inquiry on child nutritional status assessment

Description of the local method

- Discussion on local malnutrition types and forms
- Discussion on signs/symptoms
- Discussion on diagnosis process

Planning the Pilot Phase

- Selecting expert caregivers
- Planning
- Proposal of the expert cargivers to be involved in the assessment

3

Pilot survey

- Team: Expert cargivers and investigator
- Diagnosis of 10 children
- Documentation of diagnosis process

Briefing

- Presentation of pilot survey results to the mothers
- Discussion on new signs/symptoms

Data collection and analysis

- Groups discussion
- The recordings of the interviews were transcribed and then content analysed using MAXQDA software

Acknowledgement

NaviNut project is funded by the Federal Ministry of Food and Agriculture (BMEL) based on a decision of the Parliament of the Federal Republic of Germany via the Federal Office for Agriculture and Food (BLE). We would like to thank the mothers from Nikki and Banikoara.



Starting point of reliable maternal malnutrition screening system

Results


1-Category of malnutrition

Ethnicity	Malnutrition forms			
	Normal	Less Severe	Moderately Severe	Severe
Gando	DJANDOUDÔ	PEOURI GONAN KOURI/KAWÊ	PEOURI TAKINORI	«PEOURI FOROUA
Baatonu	BOUANDO	POUTANKOU	-	TIKPIROU
Fulani	DJANDOUDÔ	«DANERI	-	PEOURI

2-Diagnostic process of local method: Decision tree

Baatonu: 11 signs/symtoms

Gando: 16 signs/symtoms



Fulani: 10 signs/symtoms

Weight loss

Child still quiet

Peouri daneri

The child does

not eat well anymore

Peouri daneri

Legend

The child sleeps all the

Other disease

Other disease

5 Simillar signs/symtoms

· Oedema of cheeks and feet

Heavy

breathing

Normal

disease

Bloated

belly

- Child eating sand or clay
 - · Child not eating well
 - Weight loss
 - Pale eyes

Peouri Signs/symptoms Diagnostic result

Soft body/Edema on

cheek and feet/soft

The child does not

eat well anymore

disease

Peouri

Child eats sand or

Conclusion



Similarities between Possible use of their the methods reflecting local expertise



differences to improve each method through mutual learning



Possible integration into the current communitybased nutrition surveillance system



Chérif ISSIFOU PhD student, University of Abomey-Calavi

Corresponding author

ic.issifou@yahoo.com









